

Total Hip Arthroplasty Clinical Pathway Plan of Care

cosii	suanty for living	Date:
)	Physical Activity in conjunction with Therapy:	
•	Precautions:	

Readiness for Enhanced Therapeutic Management
R/T: impaired physical function, pain and insufficient knowledge of condition,
medications, diet, and safety secondary to total hip arthroplasty. Over the next
20 days the patient will:

Nursing	
Goals	Interventions
. Patient will remain free of skin related breakdown or infection to surgical incision.	 Monitor skin with showers and assess wound daily for s/s of infection (temp, swelling, drainage, and erythema) Provide treatments as ordered Preventable measures as indicated per care card
 Patient will remain free from pain that impacts ability to rest or participate in care. 	 Complete the pain assessment tool on admission Assess pain interfering with eating, rest and with activity Offer non-pharm interventions: Refer to pain committee and physician as needed for uncontrolled pain
Patient will remain free of preventable complications of illness such as DVT and secondary infections.	 Monitor vital signs daily and as needed Monitor lab values as ordered Observe for signs of DVT – increased swelling, erythema, calf pain Observe for signs of hip dislocation – uncontrolled pain, marked internal rotation or shortening of the surgical side
Mobility/Therapy	
Goals	Interventions
ransfers with the least amount of assistance while maintaining appropriate weight bearing precautions Patient will ambulate with appropriate assistive device for 100 feet over level surfaces with the least amount of assistance for safety. Patient will verbalize understanding of hip precautions including proper positioning of the affected extremity with less than 50% verbal cues Patient will demonstrate appropriate use of adaptive equipment for ADLs. Patient will demonstrate toileting, dressing, hygiene and grooming with the least amount of assistance	 Therapeutic exercise for Upper body Lower body Core strengthening Proprioceptive and balance training to improve safety with functional activities (weight shifting, closed kinetic chain activities) ADL training to encourage highest level of independence and safety, train on adaptive equipment. As independence is increased with ADLs, focus will shift to higher level ADL's such as bathing, cooking by discharge Gait training Modalities as needed for pain control Education regarding hip precautions, therapy goals, interventions, progress toward anticipated discharge location Lymphedema therapist to provide MLD if appropriate Transfer training with appropriate assistive device Training with use of adaptive equipment for ADLs Bed mobility on flat surface Patterned electrical nerve stimulation (PENS) if poor agonist/antagonist activity noted as needed



Total Hip Arthroplasty Clinical Pathway

PLACE- consummity for keing	Date:				
Education Needs – All disciplines					
Goals	Interventions				
 Patient will verbalize and/or demonstr understanding of education provided. 	 Provide verbal and written education Educate Patient on precautions and body mechanics with adl's and mobility Offer to have family or significant others involved in explanations and teaching sessions. This can encourage their support; reinforce the need for the Patient to assume responsibility for managing as much of care as possible Monitor and educate compliance with hip precautions, DVT prevention, safety considerations and strategies to decrease risk of falls Educate patient on care plan goals, interventions and progress toward anticipated discharge location Educate on causes of skin breakdown and wound infections Educate on DVT prevention, compliance with hip precautions, safety considerations and 				
Social Services					
Goals	Interventions				
Psychosocial factors that may affect or health will be identified	verall • Accept Patients evaluation of own strengths/limitations while working				

		•	Encourage questions and clarify misconceptions	
Dietary				
	Goals		Interventions	
	aintaining weight and voicing	•	Food alternatives/increased dietary needs per protocol/utilization of	
	g of the importance of		"Real food First". Determine and honor dietary preferences/choices.	
nutrition and	hydration for healing.	•	Weekly weight or weights as ordered	
2. Patient will m	aintain fluid balance	•	Monitor for s/s of dehydration or fluid volume overload	
Transition to Home				
	Goals		Interventions	
1. Patient will tra	ansition to home by day 21	•	All equipment and services needed are obtained prior to discharge	
with all resour	rces in place for success	•	Home health or outpatient orders in place	
Other				
	Goals		Interventions	
		•		
		•		
		•		
		•		
Staff/Title/Date:			Staff/Title/Date:	
Staff/Title/Date:	Staff/Title/Date:		Staff/Title/Date:	

Staff/Title/Date:Staff/Title/Date:Staff/Title/Date:Staff/Title/Date:Staff/Title/Date:Staff/Title/Date:Staff/Title/Date:Staff/Title/Date: